



MEDICATION GUIDE

Roweepra (row EE pra) (levetiracetam USP) tablets, for oral use

Read this Medication Guide before you start taking Roweepra and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about Roweepra?
Like other antiepileptic drugs, Roweepra (levetiracetam) may cause suicidal thoughts or actions in a very small number of people, about 1 in 500 people taking it. Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Do not stop Roweepra (levetiracetam) without first talking to a healthcare provider.

- Stopping levetiracetam suddenly can cause serious problems. Stopping a seizure medicine suddenly can cause seizures that will not stop (status epilepticus).
- Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.
- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

What is Roweepra?
Roweepra is a prescription medicine taken by mouth that is used to treat partial-onset seizures in people 1 month of age and older. Roweepra (levetiracetam) is a prescription medicine taken by mouth that is used with other medicines to treat:

- myoclonic seizures in people 12 years of age and older with juvenile myoclonic epilepsy.
- primary generalized tonic-clonic seizures in people 6 years of age and older with certain types of generalized epilepsy.

It is not known if levetiracetam is safe or effective in children under:

- 1 month of age to treat partial-onset seizures
- 12 years of age to treat myoclonic seizures

- 6 years of age to treat primary generalized tonic-clonic seizures

Before taking your medicine, make sure you have received the correct medicine. Compare the name above with the name on your bottle and the appearance of your medicine with the description of levetiracetam tablets provided below. Tell your pharmacist immediately if you think you have been given the wrong medicine.

Who should not take Roweepra (levetiracetam)?
Do not take levetiracetam tablets if you are allergic to levetiracetam.

What should I tell my healthcare provider before starting Roweepra (levetiracetam)?
Before taking levetiracetam, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had depression, mood problems or suicidal thoughts or behavior.
- have kidney problems.
- are pregnant or planning to become pregnant. It is not known if levetiracetam will harm your unborn baby. You and your healthcare provider will have to decide if you should take levetiracetam while you are pregnant. If you become pregnant while taking levetiracetam, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can enroll in this registry by calling 1-888-233-2334 or go to <http://www.aedpregnancyregistry.org>. The purpose of this registry is to collect information about the safety of levetiracetam and other antiepileptic medicine during pregnancy.
- are breastfeeding or plan to breastfeed. Levetiracetam can pass into your breast milk. It is not known if the levetiracetam that passes into your breast milk can harm your baby. Talk to your doctor about the best way to feed your baby while you receive levetiracetam tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Do not start a new medicine without first talking with your healthcare provider.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take Roweepra (levetiracetam)?

- Take Roweepra exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much levetiracetam to take and when to take it. Levetiracetam is usually taken 2 times each day.
- Your healthcare provider may change your dose. **Do not** change your dose without talking to your healthcare provider.
- Take levetiracetam with or without food.
- Swallow the tablets whole. **Do not** chew or crush tablets. Ask your healthcare provider for levetiracetam oral solution if you cannot swallow tablets.
- If you take too much levetiracetam, call your local Poison Control Center or go to the nearest emergency room right away. pharmacist for instructions on how to use the measuring device the right way.

What should I avoid while taking Roweepra (levetiracetam)?
Do not drive, operate machinery or do other dangerous activities until you know how levetiracetam tablets affect you. Levetiracetam tablets may make you dizzy or sleepy.

What are the possible side effects of Roweepra (levetiracetam)?
Levetiracetam can cause serious side effects including:

- See “What is the most important information I should know about levetiracetam?”

Call your healthcare provider right away if you have any of these symptoms:

- mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. A few people may get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs) and unusual behavior.
- extreme sleepiness, tiredness, and weakness
- allergic reactions such as swelling of the face, lips, eyes, tongue, and throat, trouble swallowing or breathing, and hives.
- a skin rash. Serious skin rashes can happen after you start taking levetiracetam. There is no way to tell if a mild rash will become a serious reaction.
- problems with muscle coordination (problems walking and moving).

The most common side effects seen in people who take levetiracetam include:

- sleepiness
- weakness
- infection
- dizziness

The most common side effects seen in children who take levetiracetam tablets include, in addition to those listed above include:

- tiredness
- acting aggressive
- decreased appetite
- nasal congestion
- irritability

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of Roweepra (levetiracetam). For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Roweepra (levetiracetam) tablets?

- Store Roweepra (levetiracetam) tablets at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature] away from heat and light.
- Keep Roweepra tablets and all medicines out of the reach of children.**

General information about Roweepra (levetiracetam).
Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Roweepra for a condition for which it was not prescribed. Do not give Roweepra to other people, even if they

have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider information about levetiracetam that is written for health professionals.

What are the ingredients of Roweepra (levetiracetam) tablets?
Roweepra tablet active ingredient: levetiracetam, USP
Inactive ingredients: colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, polyethylene glycol 400, povidone, sodium starch glycolate, talc, titanium dioxide. The 250 mg tablets contain FD&C Blue #2 Lake of indigo carmine. The 500 mg tablets contain Ferric oxide yellow. The 750 mg tablets contain Ferric oxide red and FD&C #6 lake of sunset yellow.

Roweepra (levetiracetam) tablets do not contain lactose or gluten.

Manufactured by:
TORRENT PHARMACEUTICALS LTD., INDIA.

Manufactured for:
OWP Pharmaceuticals, INC.
400 E. Diehl Road, Suite 400, Naperville, IL 60563.

8094113

Revised: September 2023

For more information, go to www.torrentpharma.com or call 1-800-912-9561.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Supersedes A/W No.:		LOCATION : -		COUNTRY : US		Roweepra (levetiracetam USP) tablets		V. No.: 01	
REMARK :		NO. OF COLORS: 1		PANTONE SHADE NOS.:		Medication Guide		Date	
SUBSTRATE :		Black		Front/ Back		Front/ Back		Signature	
Activities		Department		Pkg.Dev		8094113		Name	
Prepared By		Reviewed By		Approved By		150 x 350		Date	
Pkg.Dev		Pkg.Dev		Quality		S/S		Signature	
Font size 10 pt		Font size 10 pt		Font size 10 pt		21-09-2023		Signature	

